

A novel real-time TaqMan™ PCR assay for simultaneous detection of Neotropical fox species using noninvasive samples based on *cytochrome c oxidase subunit II*

M. Cosse¹  · F. Grattarola¹ · N. Mannise¹

Received: 6 March 2017 / Accepted: 30 June 2017 / Published online: 21 July 2017
© Mammal Research Institute, Polish Academy of Sciences, Białowieża, Poland 2017

Abstract Strategies to evaluate and monitor elusive mammal species require the development of genetic techniques and their application to unambiguous biological material for ecological and genetic studies. In order to assess *cytochrome c oxidase subunit II* gene inter- and intraspecific variations, we compared sequences from different Neotropical canids and domestic dogs. We developed a primer pair to amplify a 154-bp fragment of this gene and a species-specific multiplex TaqMan™ assay for accurate identification of two native fox species occurring in sympatry in South America, the crab-eating fox (*Cerdocyon thous*) and the pampas fox (*Lycalopex gymnocercus*). The assays can also distinguish domestic dogs (*Canis lupus familiaris*) from both wild foxes. The use of different fluorescent reporter dyes for species identification in a multiplex probe PCR-RT assay reduces labor and costs. The methodology presented in this study demonstrates an efficient approach to enable high-performance analysis and represents a reliable cost-effective tool for molecular ecology research to monitor the wild canid populations by noninvasive genetic sampling. This standardized assay will allow large-scale high-throughput analyses in a routine and reliable way.

Keywords mtDNA · *COII* gene · Molecular ecology · Canid · Biodiversity monitoring · Uruguay

Communicated by: Cino Pertoldi

✉ M. Cosse
marianacosse@gmail.com

¹ Genética de la Conservación, Departamento de Biodiversidad y Genética – IIBCE/MEC, Av. Italia 3318, 11600 Montevideo, Uruguay

Introduction

In the Neotropical Region (Procheş and Ramdhani 2012; Morrone 2014), two wild fox species are sympatric: the crab-eating fox (*Cerdocyon thous*) and the pampas fox (*Lycalopex gymnocercus*) (Fig. 1). In addition to their overlapping distribution range, they have a similar body size and share eating habits (Vieira and Port 2007). The crab-eating fox is the widest distributed canid of South America, ranging from Uruguay and northern Argentina to Paraguay, Colombia, Venezuela, and throughout Brazil, except for the Amazon basin lowlands (Berta 1982). Even though the species is listed as Least Concern for the IUCN (International Union for Conservation of Nature) (Courtenay and Maffei 2010), it is on Appendix II of CITES as a vulnerable species with regard to exploitation and trade activities (Nowak 1999). The pampas fox has a typical southern geographical distribution, occurring in eastern Bolivia, western and central Paraguay, Uruguay, north and central Argentina, and southeastern Brazil. It has also been assigned to the Least Concern category of the IUCN (Lucherini and Luengos Vidal 2008) and is included on Appendix II of CITES. The pampas fox seems to occupy only open areas including grasslands and dirt roads. Meanwhile, the crab-eating fox is found in more diverse habitats including the interior of forests as well as the edge of forests and grasslands, but more frequently in woodland areas (Lucherini et al. 2004). In addition, in natural areas, it is possible to find domestic dog (*Canis lupus familiaris*). The Uruguayan Zoonosis Commission estimates more than 146,000 dogs inhabit rural areas (Lete J., personal communication) so it is common sampling areas with dogs present. The high number of dogs present in such areas is affecting wild fauna because they predate small- and medium-sized animals such as southern tamandua (*Tamandua tetradactyla*). Transmission of diseases such as rabies and canine distemper




Fig. 1 Map of Uruguay showing the results in different sample types (control, fur, or feces) and fox species (*Cerdocyon thous*: CT; *Lycalopex gymnocercus*: LGY). *Top right*: fox species geographic distribution and sympatric area

virus may also affect wildlife (Hughes and Macdonald 2013). In Uruguay, studies on conflicts between domestic dogs and wildlife are needed to suggest accurate management activities.

Samples obtained from noninvasive methods, such as feces or shed hairs, are recognized as a valuable source of DNA in different molecular approaches for assessing genetic structure,

demography, and life history of mammals (Kohn and Wayne 1997; Beja-Pereira et al. 2009; Hausknecht et al. 2010; Ebert et al. 2012). Hence, the identification at individual or species level obtained by genetic analysis from noninvasive samples allows to monitor the presence and distribution of elusive species and to estimate their abundance applying capture-

recapture methods (Petit and Valiere 2006; Mondol et al. 2009; O'Meara et al. 2014). The first key step in fecal and other noninvasive biological sample analysis is accurately assigning species; this provides needed information to resolve continuing further molecular analyses.

More precise, fast, and inexpensive methods for identification of species are thus sought by molecular ecology and conservation genetics researchers. The amplification of small fragments (ca. 150-bp) from mitochondrial DNA (mtDNA) by polymerase chain reaction (PCR) is a powerful tool for species identification (Chaves et al. 2012; O'Meara et al. 2014; Rodriguez-Castro et al. 2017). Additionally, different TaqMan™ probes in multiplex assays is a well-documented technique to determine DNA sequence variation (Holland et al. 1991; Schoske et al. 2003; Ali et al. 2014). This methodological approach in combination with real-time PCR (PCR-RT) assays is the technique of choice for rapid, accurate, and sensitive analysis (Walker 2002) and even for low template DNA concentrations (O'Neill et al. 2013).

In this study, we report the development of a novel TaqMan™ probe PCR-RT assay based on the sequence of a fragment of the *cytochrome c oxidase subunit II (COII)* gene (mtDNA), which can be used to discriminate sympatric Neotropical Canidae species and dogs.

Materials and methods

Skin, muscle, and liver tissues from crab-eating fox (*C. thous*) ($n = 13$), pampas fox (*L. gymnocercus*) ($n = 8$), and ambiguous or non-accurate species determination canids ($n = 112$) were obtained from roadkill animals in Uruguay, collected between 1997 and 2016. Blood samples from domestic dogs ($n = 10$) were obtained from breeders. Additionally, fecal samples with a preliminary indication on canid species (ambiguous or non-accurate) were collected during surveys in Uruguay ($n = 112$) between 2009 and 2016. All tissue and fecal samples were stored in 95% ethanol at 4 °C just after collection and until DNA extraction. DNA was isolated from tissue samples following González et al. (2015) protocol. To isolate DNA from feces, we used QIAamp DNA Stool Mini Kit® (QIAGEN, Hilden, Germany) according to the manufacturer's instructions. This protocol for DNA isolation was carried out in a separate room with several safeguards against contamination including laminar flow chambers, UV lamps, and dedicated equipment and reagents which are used neither for tissue samples nor post-PCR applications. In addition, extraction negative controls were included. DNA was eluted in 100 µL H₂O for all sample types and stored at 4 °C. Quantification of DNA from tissue and fur samples was carried out in a NanoDrop™ ND-1000 UV-vis Spectrophotometer (NanoDrop Technologies, Inc., Wilmington, DE).




Fig. 2 Gene tree derived from *COII* gene based on 608-bp data sets containing sequences from 35 canids and *Phoca caspica* (PC) as external outgroup. Neighbor-Joining method (NJ) with bootstrap replicates (1000). Numbers above the branches indicate nodes supported in 90% of bootstrap replicates. LGY: *L. gymnocercus*; LV: *L. vetulus*; LGR: *L. griseus*; LC: *L. culpaeus*; LS: *L. sechurae*; CT: *C. thous*; CB: *C. brachyurus*; CLF: *C. l. familiaris*; SV: *S. venaticus*. GenBank accession number are shown; those starting with KY correspond to sequences obtained from this study

In order to assess *COII* inter- and intraspecific variation, we compared sequences from different Neotropical canids and domestic dogs deposited in GenBank (ncbi.nlm.nih.gov), using ClustalW tool in Mega6 (Tamura et al. 2013). The alignments were compared using *Canis lupus familiaris* ($n = 101$), *C. thous* ($n = 2$: AF028217, AY609148), *L. gymnocercus* ($n = 2$: AY609150, AF028225), *Lycalopex vetulus* ($n = 1$: AF028220.1), *Lycalopex culpaeus* ($n = 1$: AF028223.1), *Lycalopex griseus* ($n = 1$: AF028224.1), *Lycalopex sechurae* ($n = 2$: AF028226.1, KT448284.1), *Speothos venaticus* ($n = 3$:

Table 1 TaqMan™ probe developed. Probe IDs correspond with CLF: *C. lupus familiaris*, CT: *C. thous*, LGY: *L. gymnocercus*

TaqMan™ MGM probe ID	Probe sequence (5'–3')	Reporter	Quencher	T _m (°C)	Length (bp)
CLF FOX COX2 PROBE	C CAC AGC TTT ATA CCC ATT GTT CTT GAA A	CY5	BHQ2	66.2	29
CT FOX COX2 PROBE	AC AGC TTT ATA CCT ATT GTC CTC GAA A	HEX	BHQ1	63.7	27
LGY FOX COX2 PROBE	AC AGC TTC ATA CCC ATC GTT CTT GAA A	FAM	BHQ1	65.3	27

AF028227.1, KT448285.1, AY609154.1) and *Chrysocyon brachyurus* ($n = 2$: AY609144.1, KJ508409.1).

Tissue samples from different canid species-identified individuals were amplified using COII-F (5'-GTAA

AACATTACATGACTTTTGTC-3') and COII-R (5'-AGAG GTTAAACTCCCAGTCTT-3') (Bardeleben et al. 2005a). Each PCR contained 0.2 U Taq DNA Polymerase (Invitrogen), 1X PCR Buffer, 0.2 μ M MgCl₂, 0.4 μ M each




Fig. 3 Real-time TaqMan™ method for species identification. Probe hybridization of specimens. **a** Fam probe. **b** Hex probe. **c** Cy5 probe. **d** Allelic discrimination of haplotypes by auto analysis software 6.0 program of Rotor-Gene after amplification. **e** Cluster plots. The x-

axis and y-axis represents the signal intensities to each channel (*green*, *yellow*, and *red*). Every *dot* symbolizes the signal for each probe from one individual. *Colors* represent the assigned species: *green*: LGY; *black*: CT; *red*: CLF; *purple*: negative controls

Fig. 4 *COII* 18-bp sequences showing the specie-specific polymorphism and the synonymous substitutions observed in the *COII* binding probe region

Mitochondrial genome position (bp)	7651	7654	7660	7663	7666	7669
Fox probes position (bp)	5	8	14	17	20	23
Transitions C/T in third codon position	AGT/C	TTT/C	CCT/C	ATT/C	GTT/C	CTT/C
Aminoacid	Ser (S)	Phe (F)	Pro (P)	Ile (I)	Val (V)	Leu (L)
<i>Canis lupus familiaris</i>	C	T	C	T	C	C
<i>Cerdocyon thous</i>	C	T	T	C	T	T
<i>Lycalopex sp. Except Lycalopex sechurae</i>	C	C	C	T	C	C
<i>Lycalopex sechurae</i>	T	T	C	C	T	T
<i>Speothos venaticus</i>	C	T	T	T	T	C
<i>Chrysocyon brachyurus</i>	C	T	T	T	T	T

primer, template (≈ 90 ng DNA), and H₂O to a final volume of 15 μ L. The cycling conditions began with an initial step of 10 min at 94 °C, followed by 40 cycles of 30 s at 94 °C, 30 s at 55 °C, 90 s at 72 °C, and a final step of 20 min at 72 °C using sterile filtered pipette tips. PCR products were purified and sequenced with an ABI3130 (Applied Biosystems) at MacroGen Korea sequencing service. Sequences were aligned using ClustalW in Mega 6 (Tamura et al. 2013) including a sample of *Phoca caspica* (PC) as external outgroup. The sequence similarity among species were evaluated using the neighbor-joining (NJ) algorithm (Saitou and Nei 1987) as executed in MEGA 6 with the Kimura 2-parameter (K2P) model (Kimura 1980) of nucleotide substitution. The degree of information support for clades was assessed after bootstrap resampling of 1000 pseudo-replicates (Felsenstein 1985).

We developed a primer pair for the amplification of a short fragment of mtDNA *COII* from *Cerdocyon*, *Lycalopex*, and *Canis* species. Three TaqManTM probes were designed using intraspecies conserved sequences to detect *C. thous*, *L. gymnocercus*, and *Canis l. familiaris*. Primer Express 3.0 software from Applied Biosystems (Life Technologies) was used. The TaqManTM fluorescent probes were labeled with reporter and quencher dyes with different absorption spectra (Johansson 2006).

Novel primers and probes were validated by PCR-RT assays using DNA isolated from tissue of known-species canids ($n = 24$). Each PCR contained 1X SensiFastprobeTM PCR master mix (Bioline), 2 mM MgCl₂, 0.4 μ M each primer, 0.15 μ M each probe, template (≈ 60 ng DNA), and H₂O to a final volume of 20 μ L. The analysis was conducted using the following cycling conditions on a Rotor-Gene 6000 (Corbett Research) thermocycler: initial step of 5 min at 95 °C, followed by 40 cycles of 10 s at 95 °C and 60 s at 62 °C. Fluorescence acquisition was determined during the extension step on the green channel (470 nm excitation, 510 nm detection), on the yellow channel (530 nm excitation, 550 nm detection), and on the red channel (625 nm excitation, 660 nm detection) to detect the binding of the probe to the target sequences of *L. gymnocercus*, *C. thous*, or *C. lupus familiaris*, respectively. Positive and negative

controls were included in all PCR reactions. The acquisition curves analyses were performed using Rotor-GeneTM 6000 software v.1.7 and the algorithm provided. Additionally, the efficiency of the TaqManTM probes developed for species identification were assessed using DNA obtained from noninvasive samples like feces ($n = 112$). In these cases were conducted at least two replicates for each PCR with 1 and 2 μ L DNA volume. From seven roadkilled foxes, we collected tissue and feces present in the rectum. We conducted real-time PCR TaqManTM probe on both types of samples collected from the same animal. Furthermore, in four feces samples, we performed not only the probe experiment but also sequenced the *COII* probe region.

Results

We obtained 17 sequences from *COII* mtDNA [*C. thous* ($n = 8$), *L. gymnocercus* ($n = 5$), and *C. l. familiaris* ($n = 4$)]. The fragments generated were 608-bp long. For the *COII* analysis, the alignment was constructed with sequences available from GenBank and those obtained in this work, summing 35 sequences (608-bp) that included eight Neotropical canid species and domestic dogs. Except *L. gymnocercus*, all species represented by more than one sequence formed monophyletic groups ($\geq 99\%$ BS) (Fig. 2). Conserved flanking region sequences for PCR primer design were identified. We also identified sequence variation that could be used to distinguish canid species present in Uruguay: *C. l. familiaris*, *C. thous*, and *L. gymnocercus* as well as *S. venaticus*, *C. brachyurus*, and *L. sechurae*.

The primer pair designed (FOX154F: CGC TAT YCC AGG ACG ACT A and FOX154R: GGC TGA TCA GGT TTC AAA G) successfully amplified a 154-bp fragment of *COII* mtDNA and the binding site for TaqManTM probes designed (Table 1) for *C. thous*, *L. gymnocercus*, and *C. l. familiaris* were also unequivocally validated by assaying all control samples, i.e., samples of crab-eating foxes ($n = 14$), pampas foxes ($n = 10$), and dogs ($n = 4$).

From 112 roadkilled samples, 95 were correctly identified and it represents that 84.8% of the experiments were successful. In 71.4% of cases, the result were obtained in the first experiment, while in 13.4%, it was achieved in the second one. The 15.2% of PCR did not amplify in these kind of samples. From roadkilled samples, 64 were crab-eating foxes (*C. thous*) and 31 were pampas foxes (*L. gymnocercus*). Only in one case we detected a misidentification in a sample that was primary identified as *L. gymnocercus* and resulting in *C. thous* from two independent PCRs. In addition, from 112 feces samples we identified 26 as crab-eating foxes (*C. thous*) and 40 as pampas foxes (*L. gymnocercus*), i.e., 58.9% of the feces samples could be used for identification at the species level (Fig. 3). Feces showed 48.2% of success on the first PCR amplifications and only 10.7% of samples were identified after replicates. The accuracy of this technique with noninvasive samples like feces was confirmed from the same and stable pattern of Taqman™ probes with fur and rectal samples from the same individuals in seven evaluated cases. Finally, the sequenced *COII* fragment from feces samples showed concordance with the probe experiments.

Discussion

COII gene (partial region of 608-bp) was suitable for Neotropical fox species identification with high bootstrap values for each clade, except for *L. gymnocercus*, which is not represented as a monophyletic clade (Fig. 2). This pattern in *L. gymnocercus* was observed by Tchaicka et al. (2016) with mtDNA control region and these authors suggested an inter-species hybridization process or a secondary admixture. Nevertheless, the *COII* analyzed region (608-bp) shows homoplasy and the tree performed with NJ method based in this region seem not to reflect the phylogenetic relation of the species (Pamilo and Nei 1988; Wayne et al. 1997; Bardeleben et al. 2005b; Bardeleben et al. 2005a; Galimberti et al. 2015).

For noninvasive DNA sampling, mtDNA sequences tend to result in better amplification success (70–90%) against nuclear markers (Broquet et al. 2007). The novel primer pair designed in this study could be used for PCR amplification of a 154-bp *COII* gene fragment in the Neotropical canid species and the domestic dog. The length of the PCR product was suitable for amplification of low-quality and low-quantity DNA, given that amplification success of fecal DNA from carnivore species was 71.9%.

The TaqMan™ probe binding region (18-bp) has synonymous substitutions with a polymorphism level that allows distinguishing Neotropical canid species in Uruguay. Additionally, this 18-bp region has potential for being useful to identify several other Neotropical canid species (Fig. 4). Nevertheless, the *L. gymnocercus* TaqMan™ probe binds to

all the species of the genus (*L. culpaeus*, *L. griseus*, *L. vetulus*) except for *L. sechurae*. The polymorphism for *Lycalopex fulvipes* was not analyzed because there were no sequences available in GenBank.

The use of different fluorescent reporter dyes for species identification in a multiplex probe PCR-RT assay reduces labor and costs. The methodology presented in this study demonstrates an efficient approach to facilitate high-throughput analysis and critical evaluation of *C. thous*, *L. gymnocercus*, and dog samples in a single analysis. The Scatter Analysis Graph with the software provided by Rotor-Gene Q (V.2.3.1) allows an easy taxonomic determination (Fig. 3e). This standardized assay will allow large-scale high-throughput analyses in a routine and reliable way.

Acknowledgements The research was supported by *Programa de Desarrollo de las Ciencias Básicas* (PEDECIBA) and grant from *Agencia Nacional de Investigación e Innovación* (ANII) (FCE_2_20011_1_5700). The authors wish to thank the *Monitoreo participativo de fauna en Paso Centurión EFI-UdelaR* team, and Cabo Polonio forest ranger team (SNAP) and to acknowledge sample contribution from C. Máspoli, C. Aristimuño, C. Prigioni, M. Baptista, A. Saralegi, O. Lussich, N. Hernandez and C. Pérez, and José M. Venzal.

References

- Ali ME, Razzak MA, Hamid SBA (2014) Multiplex PCR in species authentication: probability and prospects—a review. *Food Anal Methods* 7(10):1933–1949
- Bardeleben C, Moore RL, Wayne RK (2005a) Isolation and molecular evolution of the selenocysteine tRNA (Cf TRSP) and RNase P RNA (Cf RPPH1) genes in the dog family, Canidae. *Mol Biol Evol* 22(2): 347–359
- Bardeleben C, Moore RL, Wayne RK (2005b) A molecular phylogeny of the Canidae based on six nuclear loci. *Mol Phylogenet Evol* 37(3): 815–831
- Beja-Pereira A, Oliveira R, Alves PC, Schwartz MK, Luikart G (2009) Advancing ecological understandings through technological transformations in noninvasive genetics. *Mol Ecol Resour* 9(5):1279–1301
- Berta A (1982) *Cerdocyon thous*. *Mamm Species* 186:1–4
- Broquet T, Ménard N, Petit E (2007) Noninvasive population genetics: a review of sample source, diet, fragment length and microsatellite motif effects on amplification success and genotyping error rates. *Conserv Genet* 8(1):249–260
- Chaves PB, Graeff VG, Lion MB, Oliveira LR, Eizirik E (2012) DNA barcoding meets molecular scatology: short mtDNA sequences for standardized species assignment of carnivore noninvasive samples. *Mol Ecol Resour* 12(1):18–35
- Courtenay O, Maffei L (2010) *Cerdocyon thous*. In: Sillero-Zubiri C, Hoffmann M, Macdonald DW (eds) *Canids: foxes, wolves, jackals and dogs. Status survey and conservation action plan: IUCN/SSC Canid Specialist Group, IUCN, Gland. IUCN, Cambridge*, pp 32–38
- Ebert C, Knauer F, Spielberger B, Thiele B, Hohmann U (2012) Estimating wild boar *Sus scrofa* population size using faecal DNA and capture-recapture modelling. *Wildl Biol* 18(2):142–152
- Felsenstein J (1985) Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* 39(4):783–791

- Galimberti A, Sandionigi A, Bruno A, Bellati A, Casiraghi M (2015) DNA barcoding in mammals: what's new and where next? *Hystrix* 26(1):13–24
- González S, Cosse M, del Rosario FM, Emmons L, Vynne C, Duarte JMB, Beccacesi MD, Maldonado JE (2015) Population structure of mtDNA variation due to Pleistocene fluctuations in the South American Maned Wolf (*Chrysocyon brachyurus*, Illiger, 1815): management units for conservation. *J Hered* 106(S1):459–468
- Hausknecht R, Bayerl H, Gula R, Kuehn R (2010) Application of quantitative real-time polymerase chain reaction for noninvasive genetic monitoring. *J Wildl Manag* 74(8):1904–1910
- Holland PM, Abramson RD, Watson R, Gelfand DH (1991) Detection of specific polymerase chain reaction product by utilizing the 5'—3'exonuclease activity of *Thermus aquaticus* DNA polymerase. *P Natl Acad Sci USA* 88(16):7276–7280
- Hughes J, Macdonald DW (2013) A review of the interactions between free-roaming domestic dogs and wildlife. *Biol Conserv* 157:341–351
- Johansson MK (2006) Choosing reporter-quencher pairs for efficient quenching through formation of intramolecular dimers. In: *Fluorescent Energy Transfer Nucleic Acid Probes*. Springer, pp 17–29
- Kimura M (1980) A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *J Mol Evol* 16(2):111–120
- Kohn MH, Wayne RK (1997) Facts from feces revisited. *Trends Ecol Evol* 12(6):223–227
- Lucherini M, Luengos Vidal EM (2008) *Lycalopex gymnocercus* (Carnivora: Canidae). *Mamm Species* 1–9
- Lucherini M, Pessino M, Farias AA (2004) Pampas fox (*Pseudalopex gymnocercus*). In: Sillero-Zubiri C, Hoffmann M, Macdonald DW (eds) *Canids: foxes, wolves, jackals and dogs: status survey and conservation action plan*. Survey and conservation action plan, vol Status. IUCN/SSC Canid Specialist Group, Gland, pp 63–68
- Mondol S, Ullas Karanth K, Samba Kumar N, Gopaldaswamy AM, Andheria A, Ramakrishnan U (2009) Evaluation of non-invasive genetic sampling methods for estimating tiger population size. *Biol Conserv* 142(10):2350–2360
- Morrone JJ (2014) Biogeographical regionalization of the Neotropical region. *Zootaxa* 3782(1):1–110
- Nowak NM (1999) *Walker's mammals of the world*, vol 1. JHU Press, Baltimore
- O'Meara DB, Sheehy E, Turner PD, O'Mahony D, Harrington AP, Denman H, Lawton C, MacPherson J, O'Reilly C (2014) Non-invasive multi-species monitoring: real-time PCR detection of small mammal and squirrel prey DNA in pine marten (*Martes martes*) scats. *Acta Theriol* 59(1):111–117
- O'Neill D, Turner PD, O'Meara DB, Chadwick EA, Coffey L, O'Reilly C (2013) Development of novel real-time TaqMan® PCR assays for the species and sex identification of otter (*Lutra lutra*) and their application to noninvasive genetic monitoring. *Mol Ecol Resour* 13(5):877–883
- Pamilo P, Nei M (1988) Relationships between gene trees and species trees. *Mol Biol Evol* 5(5):568–583
- Petit E, Valiere N (2006) Estimating population size with noninvasive capture-mark-recapture data. *Conserv Biol* 20(4):1062–1073
- Procheş Ş, Ramdhani S (2012) The world's zoogeographical regions confirmed by cross-taxon analyses. *Bioscience* 62(3):260–270
- Rodriguez-Castro KG, Ciocheti G, Ribeiro JW, Ribeiro MC, Galetti PM (2017) Using DNA barcode to relate landscape attributes to small vertebrate roadkill. *Biodivers Conserv*:1–18
- Saitou N, Nei M (1987) The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* 4(4):406
- Schoske R, Vallone PM, Ruitberg CM, Butler JM (2003) Multiplex PCR design strategy used for the simultaneous amplification of 10 Y chromosome short tandem repeat (STR) loci. *Anal Bioanal Chem* 375(3):333–343
- Tamura K, Stecher G, Peterson D, Filipinski A, Kumar S (2013) MEGA6: molecular evolutionary genetics analysis version 6.0. *Mol Biol Evol* 30:2725–2729
- Tchaicka L, Freitas TRO, Bager A, Vidal SL, Lucherini M, Iriarte A, Novaro A, Geffen E, Garcez FS, Johnson WE (2016) Molecular assessment of the phylogeny and biogeography of a recently diversified endemic group of South American canids (Mammalia: Carnivora: Canidae). *Genet Mol Biol* 39(3):442–451
- Vieira E, Port D (2007) Niche overlap and resource partitioning between two sympatric fox species in southern Brazil. *J Zool* 272(1):57–63
- Walker NJ (2002) A technique whose time has come. *Science* 296(5567):557
- Wayne RK, Geffen E, Girman DJ, Koepfli KP, Lau LM, Marshall CR (1997) Molecular systematics of the Canidae. *Syst Biol* 46(4):622–653